arXiv:1607.07578v3 [math.OA] 22 Apr 2017

Reduction to dimension two of local spectrum for AH algebra with ideal property

Chunlan Jiang

Abstract: A C^*-algebra A has ideal property if any ideal I of A is generated as a closed two sided ideal by the projections inside the ideal. Suppose that the limit C^*-algebra A of inductive limit of direct sums of matrix algebras over spaces with uniformly bounded dimension has ideal property. In this paper we will prove that A can be written as an inductive limit of certain very special subhomogeneous algebras, namely, direct sum of dimension drop interval algebras and matrix algebras over 2-dimensional spaces with torsion H^2 groups.

§1. Introduction

An AH algebra is a nuclear C^*-algebra of the form $A = \varinjlim(A_n, \phi_{n,m})$ with $A_n = \bigoplus_{i=1}^{t_n} P_{n,i}M_{[n,i]}(C(X_{n,i}))P_{n,i}$, where $X_{n,i}$ are compact metric spaces, t_n, $[n,i]$ are positive integers, $M_{[n,i]}(C(X_{n,i}))$ are algebras of $[n,i]\times[n,i]$ matrices with entries in $C(X_{n,i})$—the algebra of complex - valued functions on $X_{n,i}$, and finally $P_{n,i} \in M_{[n,i]}(C(X_{n,i}))$ are projections (see [Bla]). If we further assume that $\sup_{n,i} \dim(X_{n,i}) < +\infty$ and A has ideal property—each ideal I of A is generated by the projections inside the ideal, then it is proved in [GJLP1-2] that A can be written as inductive limit of $B_n = \bigoplus_{i=1}^{s_n} P'_{n,i}M_{[n,i]'}(C(Y_{n,i}))P'_{n,i}$.

In this paper, we will further reduce the dimension of local spectra (that is the spectra of A_n or B_n above) to 2 (instead of 3). Namely, the above A can be written as inductive limit of direct sum of matrix algebras over the $\{pt\}, [0,1], S^1, T_{II,k}$ (no $T_{III,k}$ and S^2) and $M_l(I_k)$, where I_k is dimension drop interval algebra,

$$I_k = \{f \in C([0,1], M_k(\mathbb{C})), f(0) = \lambda\mathbf{1}_k, f(1) = \mu\mathbf{1}_k, \lambda, \mu \in \mathbb{C}\}.$$

In this paper, we will also call $\bigoplus_{i=1}^{s} M_{l_i}(I_{k_i})$ a dimension drop-algebra.
This result unifies the theorems of [DG] and [EGS] (for the rank zero case) and [Li4] (for the simple case). Note that Li′s reduction theorem was not used in the classification of simple AH algebra, and Li′s proof depends on the classification of simple AH algebra (see [Li4] and [EGL]). For our case, the reduction theorem is an important step toward the classification (see [GJL]). The proof is more difficult than Li′s case. For example, in the case of AH

algebra with ideal property, one can not remove the space S^2 without introduce $M_l(I_k)$ (for the simple case, the space S^2 is removed from the list of spaces in [EGL] without introduced dimension drop algebras). Another point is that, in the simple AH algebras, one can assume each partial map $\phi^{i,j}_{n,m}$ is injective, but in AH algebras with ideal property, we can not make such assumption. For the classification of real rank zero AH algebras, we refer to the readers [Ell1], [EG1-2], [D1-2], [G1-4] and [DG]. For the classification simple AH algebra, we refer to the readers [Ell2-3], [Li1-3], [G5] and [EGL1-2].

The paper is organized as follows.
In section 2, we will do some necessary preparation. In section 3, we will prove our main theorem.

§2. Preparation

We will adopt all the notation from section 2 of [GJLP2]. For example we refer the reader to [GJLP2] for the concepts of $G-\delta$ multiplicative maps (see definition 2.2 there), spectral variation $SPV(\phi)$ of a homomorphism ϕ (see 2.12 there) weak variation $\omega(F)$ of a finite set $F \subset QM_N(C(X))Q$ (see 2.16 there).
As in 2.17 of [GJLP2], we shall use $\bullet$ to denote any possible integer.

2.1. In this article, without lose of generality we will assume the AH algebras A are inductive limit of

$$A = \lim(A_n = \bigoplus_{i=1}^{t_n} M_{[n,i]}(C(X_{n,i})), \phi_{n,m}),$$

where $X_{n,i}$ are the spaces of $\{pt\}, [0,1], S^1, T_{II,k}, T_{III,k}$, and S^2. (Note by the main theorem of [GJLP2], all AH algebras with ideal property and with no dimension growth are corner subalgebras of the above form (also see 2.7 of [GJLP2]).)

2.2. Recall a projection $P \in M_k(C(X))$ is called a trivial projection if it is unitarily equivalent to $\begin{pmatrix} \mathbf{1}_{k_1} & 0 \\ 0 & 0 \end{pmatrix}$ for $k_1 = rank(P)$. If P is a trival projection and $rank(P) = k_1$, then

$$PM_k(C(X))P \cong M_{k_1}(C(X)).$$

2.3. Let X be a connected finite simplicial complex, $A = M_k(C(X))$, a unital $*$ homomorphism $\phi : A \longrightarrow M_l(A)$ is called a (unital) simple embedding if it is homotopic to the homomorphism $id \oplus \lambda$, when $\lambda : A \longrightarrow M_{l-1}(A)$ is defined by

$$\lambda(f) = diag(\underbrace{f(x_0), f(x_0), \cdots, f(x_0)}_{l-1}),$$

for a fixed base point $x_0 \in X$.

The following two lemmas are special cases of Lemma 2.15 of [EGS] (also see 2.12 of [EGS]).

Lemma 2.4. (c.f 2.12 or case 2 of 2.15 in [EGS]). For any finite set $F \subset A = M_n(C(T_{III,k}))$ and $\varepsilon > 0$, there is a unital simple embedding $\phi : A \longrightarrow M_l(A)$ (for l large enough) and a C^*-algebra $B \subset A$, which is a direct sum of dimension drop algebras and a finite dimensional C^*-algebra such that

$$dist(\phi(f), B) < \varepsilon, \qquad \forall\ f \in F.$$

Lemma 2.5.(see case 1 of 2.15 in [EGS]) For any finite set $F \subset M_n(C(S^2))$ and $\varepsilon > 0$, there is a unital simple embedding $\phi : A \longrightarrow M_l(A)$ (for l large enough) and a C^*-algebra $B \subset A$, which is a finite dimensional C^*-algebra such that

$$dist(\phi(f), B) < \varepsilon, \qquad \forall\ f \in F.$$

The following lemma is well known:

Lemma 2.6.(see [G5,4.40]) For any C^*-algebra A and finite set $F \subset A$, $\varepsilon > 0$, there is a finite set $G \subset A$ and $\eta > 0$ such that if $\phi : A \longrightarrow B$ is a homomorphism and $\psi : A \longrightarrow B$ is a completely positive linear map, satisfing the following

$$\|\phi(g) - \psi(g)\| < \eta, \qquad \forall\ g \in G,$$

then ψ is the $F - \varepsilon$ multiplicative.

Lemma 2.7. Let $A = M_n(C(T_{III,k}))$ or $M_n(C(S^2))$. And let a finite set $F \subset A$ and $\varepsilon > 0$, there is a diagram



with the following conditions.
(1) ϕ is a simple embedding,
(2) if $A = M_n(C(S^2))$, then B is a finite dimensional C^*-algebra, and if $A = M_n(C(T_{III,k}))$, then B is a direct sum of dimension drop C^*-algebras and a finite dimensional C^*-algebra, and ι is an inclusion,
(3) $\|\iota \circ \beta(f) - \phi(f)\| < \varepsilon$, $\forall f \in F$, and β is $F - \varepsilon$ multiplicative.

Proof. Let G and η be as Lemma 2.6 for F and ε. Apply Lemma 2.4 or Lemma 2.5 to A, $F \cup G \subset A$ and $\frac{1}{3}\min(\varepsilon, \eta)$. One can find a unital simple embedding $\phi : A \longrightarrow M_l(A)$, and

an sub-C^*-algebra $B \subset M_l(A)$ as required in condition (2) such that

$$dist(\phi(f), B) < \frac{1}{3}\min(\varepsilon, \eta), \qquad \forall\ f \in F.$$

Choose a finite $\widetilde{F} \subset B$ such that

$$dist(\phi(f), \widetilde{F}) < \frac{1}{3}\min(\varepsilon, \eta), \qquad \forall\ f \in F.$$

Since B is a nuclear C^*-algebra, there are two completely positive linear maps

$$\lambda_1 : B \longrightarrow M_N(\mathbb{C}) \quad \text{and} \quad \lambda_2 : M_N(\mathbb{C}) \longrightarrow B,$$

such that

$$\|\lambda_2 \circ \lambda_1(g) - g\| < \frac{1}{3}\min(\varepsilon, \eta), \quad \forall g \in \widetilde{F}$$

Using Arveson extension theorem, one can extend $\lambda_1 : B \longrightarrow M_N(\mathbb{C})$ to a map $\beta_1 : M_l(A) \longrightarrow M_N(\mathbb{C})$. Then it is a straight forward to prove that

$$\beta = \lambda_2 \circ \beta_1 \circ \phi : A \longrightarrow B,$$

is as desired. □

The following is modification of Theorem 3.8 of [GJLP2].

Proposition 2.8. Let $\lim_{n\to\infty}(A_n = \bigoplus_{i=1}^{t_n} M_{[n,i]}(C(X_{n,i})), \phi_{n,m})$ be AH inductive limit with ideal property, with $X_{n,i}$ being one of $\{pt\}, [0,1], S^1, T_{II,k}, T_{III,k}$, or S^2. Let $B = \bigoplus_{i=1}^{s} B^i$, where $B^i = M_{l_i}(C(Y_i))$, with Y_i being $\{pt\}, [0,1], S^1$, or $T_{II,k}$, (no $T_{III,k}$ or S^2) or $B^i = M_{l_i}(I_{k_i})$—dimension drop C^*-algebras. Suppose that

$$\widetilde{G}(= \oplus\widetilde{G^i}) \subset G(= \oplus G^i) \subset B(= \oplus B^i),$$

is a finite set, ε_1 is a positive number with $\omega(\widetilde{G^i}) < \varepsilon_1$, if $Y_i = T_{II,k}$, and L is any positive integer. Let $\alpha : B \longrightarrow A_n$ be any homomorphism. Denote

$$\alpha(\mathbf{1}_B) := R(= \oplus R^i) \in A_n(= \oplus A_n^i).$$

Let $F \subset RA_nR$ be any finite set and $\varepsilon < \varepsilon_1$ be any positive number. It follows that there are A_m, and mutually orthogonal projections $Q_0, Q_1, Q_2 \in A_m$ with

$$\phi_{n,m}(R) = Q_0 + Q_1 + Q_2,$$

a unital map $\theta_0 \in Map(RA_nR, Q_0A_mQ_0)_1$, two unital homomorphisms $\theta_1 \in Hom(RA_nR, Q_1A_mQ_1)_1$, $\xi \in Hom(RA_nR, Q_2A_mQ_2)_1$ such that
(1) $\| \phi_{n,m}(f) - (\theta_0(f) \oplus \theta_1(f) \oplus \xi(f)) \| < \varepsilon$, for all $f \in F$,
(2) there is a unital homomorphism

$$\alpha_1 : B \longrightarrow (Q_0 + Q_1)A_m(Q_0 + Q_1),$$

such that

$$\| \alpha_1(g) - (\theta_0 + \theta_1) \circ \alpha(g) \| < 3\varepsilon_1 \quad \forall g \in \widetilde{G}_i, \qquad \text{if } B^i \text{ is of form } M_\bullet(T_{II,k})$$

and

$$\| \alpha_1(g) - (\theta_0 + \theta_1) \circ \alpha(g) \| < \varepsilon, \quad \forall g \in G^i, \qquad \text{if } B^i \text{ is not of form } M_\bullet(T_{II,k}).$$

(3) θ_0 is $F - \varepsilon$ multiplicative and θ_1 satisfies that

$$\theta_1^{i,j}([e]) \geqslant L \cdot [\theta_0^{i,j}(R^i)].$$

(4) ξ factors through a C^*-algebra C—a direct sum of matrix algebras over C[0,1] as

$$\xi : RA_nR \xrightarrow{\xi_1} C \xrightarrow{\xi_2} Q_2A_mQ_2.$$

Proposition 2.9. Let $\lim_{n\to\infty}(A_n = \bigoplus_{i=1}^{t_n} M_{[n,i]}(C(X_{n,i})), \phi_{n,m})$ be AH inductive limit with ideal property, with $X_{n,i}$ being one of $\{pt\}, [0,1], S^1, T_{II,k}, T_{III,k}$ or S^2. Let $B = \bigoplus_{i=1}^{s} B^i$, where $B^i = M_{l_i}(C(Y_i))$, with Y_i being $\{pt\}, [0,1], S^1$ or $T_{II,k}$, (no $T_{III,k}$ or S^2) or $B^i = M_{l_i}(I_{k_i})$—dimension drop C^*-algebras. Suppose that

$$\widetilde{G}(= \oplus\widetilde{G}^i) \subset G(= \oplus G^i) \subset B(= \oplus B^i),$$

is a finite set, ε_1 is a positive number with $\omega(\widetilde{G}^i) < \varepsilon_1$, if $Y_i = T_{II,k}$, and $L > 0$ is any positive integer. Let $\alpha : B \longrightarrow A_n$ be any homomorphism. Let $F \subset A_n$ be any finite set and $\varepsilon < \varepsilon_1$ be any positive number. It follows that there are A_m and mutually orthogonal projections $P, Q \in A_m$ with $\phi_{n,m}(\mathbf{1}_{A_n}) = P + Q$, a unital map $\theta \in Map(A_n, PA_mP)_1$, and a unital homomorphism $\xi \in Hom(A_n, QA_mQ)_1$ such that
(1) $\| \phi_{n,m}(f) - (\theta(f) \oplus \xi(f)) \| < \varepsilon$, for all $f \in F$,
(2) there is a homomorphism $\alpha_1 : B \longrightarrow PA_mP$ such that

$$\| \alpha_1^{i,j}(g) - (\theta \circ \alpha)^{i,j}(g) \| < 3\varepsilon_1 \quad \forall g \in \widetilde{G}^i, \qquad \text{if } B^i \text{ is of form } M_\bullet(C(T_{II,k})),$$

$$\| \alpha_1^{i,j}(g) - (\theta \circ \alpha)^{i,j}(g) \| < \varepsilon \quad \forall g \in G^i, \qquad \text{if } B^i \text{ is not of form } M_\bullet(C(T_{II,k})).$$

(3) $\omega(\theta(F)) < \varepsilon$ and θ is $F - \varepsilon$ multiplicative.
(4) ξ factors through a C^*-algebra C—a direct sum of matrix algebras over $C[0,1]$ or $\mathbb{C}$ as

$$\xi : A_n \xrightarrow{\xi_1} C \xrightarrow{\xi_2} QA_mQ.$$

The proof is similar to Proposition 2.8 and is omitted.

2.10. Let $\alpha : \mathbb{Z} \longrightarrow \mathbb{Z}/k_1\mathbb{Z}$ be the group homomorphism defined by $\alpha(\mathbf{1}) = [\mathbf{1}]$, where the right hand side is the equivalent class $[\mathbf{1}]$ of $\mathbf{1}$ in $\mathbb{Z}/k_1\mathbb{Z}$. Then it is well known from homological algebra that for the group $\mathbb{Z}/k\mathbb{Z}$, α induces a surjective map

$$\alpha_* : Ext(\mathbb{Z}/k\mathbb{Z}, \mathbb{Z})(= \mathbb{Z}/k\mathbb{Z}) \longrightarrow Ext(\mathbb{Z}/k\mathbb{Z}, \mathbb{Z}/k_1\mathbb{Z})(= \mathbb{Z}/(k, k_1)\mathbb{Z}),$$

where (k, k_1) is the greatest common factor of k and k_1.
Recall, as in [DN], for two connected finite simplicial complexes X and Y, we use $kk(Y, X)$ to denote the group of equivalent classes of homomorphisms from $C_0(X\backslash\{pt\})$ to $C_0(Y\backslash\{pt\})\otimes \mathcal{K}(H)$. Please see [DN] for details.

Lemma 2.11. (a) Any unital homomorphism

$$\phi : C(T_{II,k}) \longrightarrow M_\bullet(C(T_{III,k_1})),$$

is homotopy equivalent to unital homomorphism ψ factor as

$$C(T_{II,k}) \xrightarrow{\psi_1} C(S^1) \xrightarrow{\psi_2} M_\bullet(C(T_{III,k_1})).$$

(b) Any unital homomorphism $\phi : C(T_{II,k}) \longrightarrow PM_\bullet(C(S^2))P$ is homotopy equivalent to unital homomorphism ψ factor as

$$C(T_{II,k}) \xrightarrow{\psi_1} \mathbb{C} \xrightarrow{\psi_2} PM_\bullet(C(S^2))P.$$

Proof. Part (b) is well known (see chapter 3 of [EG2]). To prove part (a), we note that

$$KK(C_0(S^1\backslash\{1\}), C_0(T_{III,k_1}\backslash\{x_1\}) = kk(T_{III,k_1}, S^1) = \mathbb{Z}/k_1\mathbb{Z} = Hom(\mathbb{Z}, \mathbb{Z}/k_1\mathbb{Z})),$$

where $x_1 \in T_{III,k_1}$ is a base point. The map $\alpha : \mathbb{Z} \longrightarrow \mathbb{Z}/k_1\mathbb{Z}$ in 2.10 can be induced by a homomorphism: $\psi_2 : C(S^1) \longrightarrow M_\bullet(C(T_{III,k}))$.
Let

$$[\phi] \in kk(T_{III,k_1}\ T_{II,k}) = Ext(K_0(C_0(T_{II,k}\backslash\{x_0\})), K_1(C_0(T_{III,k}))),$$

be the element induced by homomorphism ϕ, where $\{x_0\}$ is the base point. By 2.10

$$[\phi] = \beta \times [\psi_2],\ for\ \beta \in kk(S^1, T_{II,k}) = Ext(K_0(C(T_{II,k}\backslash\{x_0\})), K_1(C(S^1))),$$

on the other hand β can be realized by unital homomorphism

$$\psi_1 : C(T_{II,k}) \longrightarrow C(S^1).$$

(see section 3 of [EG2]).

□

The following result is modification of Theorem 3.12 of [GJLP2].

Theorem 2.12. Let $B_1 = \bigoplus_{i=1}^s B_1^i$, each B^i is either matrix algebras over $\{pt\}, [0, 1], S^1$ or $\{T_{II,k}\}_{k=2}^\infty$ or a dimension drop algebra. Let $\varepsilon_1 > 0$ and let

$$\widetilde{G}_1(= \oplus\widetilde{G}_1^i) \subset G_1(= \oplus G_1^i) \subset B_1(= \oplus B_1^i),$$

be a finite set with $\omega(G_1^i) < \varepsilon_1$ for $B_1^i = M_\bullet(C(T_{II,k}))$.

Let $A = M_N(C(X))$, where X is one of form $\{pt\}, [0,1], S^1, \{T_{II,k}\}_{k=2}^{\infty}, \{T_{III,k}\}_{k=2}^{\infty}$ or S^2. Let $\alpha_1 : B_1 \longrightarrow A$ be a homomorphism. Let $F_1 \subset A$ be a finite set, and let $\varepsilon(< \varepsilon_1)$ and δ be any positive number. Then there exists a diagram.



where $A^{'} = M_K(A)$, B_2 is described as below. B_2 is a direct sum of finite dimensional C^*-algebra and dimension drop algebra if $X = T_{III,k}$. B_2 is a finite dimensional algebra if $X = S^2$. And $B_2 = M_\bullet(A)$ if $X = \{pt\}, [0,1], S^1, T_{II,k}$. Furthermore the diagram satisfies the following conditions
(1) ψ is a homomorphism, α_2 is a unital injective homomorphism and ϕ is a unital simple embedding;
(2) $\beta \in Map(A, B_2)_1$ is $F_1 - \delta$ multiplicative;
(3) if B_1^i is of form $M_\bullet(C(T_{II,k}))$, then

$$\| \psi(g) - \beta \circ \alpha_1(g) \| < 10\varepsilon_1, \quad \forall g \in \widetilde{G}_1^i;$$

and if B_1^i is not of form $M_\bullet(C(T_{II,k}))$, then

$$\| \psi(g) - \beta \circ \alpha_1(g) \| < \varepsilon, \quad \forall g \in G_1^i;$$

(4) If $X = T_{II,k}$, then $\omega(\beta(F_1) \cup \psi(G_1)) < \varepsilon$.
(Note that, we only required that the weak variation of finite set in $M_\bullet(C(T_{II,k})$ to be small. In particular, we do not need to introduce the concept of weak variation for a finite subset of a dimension drop algebra.)

Proof. . For $X = T_{II,k}, \{pt\}, [0,1]$ or S^1, one can choose $B_2 = M_K(A) = A^{'}$ and homomorphism $\phi = \beta : A \longrightarrow B_2$ being simple embedding such that

$$\omega(\beta(F_1) \cup \alpha_1(G_1))) < \varepsilon.$$

This can be done by choosing K large enough. Choose $\psi = \beta \circ \alpha_1$, and $\alpha_2 = id : B_2 \longrightarrow A^{'}$. For the case $X = T_{III,k}$, or S^2, the requirement (4) is an empty requirement.

We will deal with each block of B_1 separately. For the block B_1^i other than $M_\bullet(C(T_{II,k})$, the construction can be done easily by using Lemma 2.7, since B_1^i is stably generated, which implies that any sufficiently multiplicative map from B_1^i is close to a homomorphism. So we assume that $B_1^i = M_\bullet(C(T_{II,k})$. Recall we already assume A is of form $M_\bullet(C(T_{III,k}))$ or $M_\bullet(C(S^2)$. By Lemma 2.11, the homomorphism $\alpha_1 : B_1^i \longrightarrow A$ is a homotopy to $\alpha^{'} : B_1^i \longrightarrow$

A with $\alpha^{'}(\mathbf{1}_{B_1^i}) = \alpha_1(\mathbf{1}_{B_1^i})$ and $\alpha^{'}$ factor as

$$B_1^i \xrightarrow{\xi_1} C \xrightarrow{\xi_2} A,$$

where C is a finite dimensional C^*-algebra for the case $X = S^2$ or $C = M_\bullet(C(S^1)$ for the case $X = T_{III,k}$ (note that $B_1^i = M_\bullet(C(T_{II,k}))$. Since C is stably generated, there is a finite set $E_1 \subset A$ and $\delta_1 > 0$ such that if a complete positive map $\beta : A \longrightarrow D$ (for any C^*-algebra D) is $E_1 - \delta_1$ multiplicative, then the map $\beta \circ \xi_2 : C \longrightarrow D$ can be perturbed to a homomorphism

$$\widetilde{\xi} : C \longrightarrow D$$

such that

$$\|\widetilde{\xi}(g) - \beta \circ \xi_2(g)\| < \varepsilon_1, \ \forall \ g \in \xi_1(\widetilde{G}_1^i).$$

Apply Theorem 1.6.9 of [G5] to two homotopic homomorphism

$$\alpha_1, \alpha^{'} : B_1^i \longrightarrow A,$$

and $G_1^i \subset B_1^i$ which is approximately constant to within ε_1, to obtain a finite set $E_2 \subset A$, $\delta_2 > 0$ and positive integer $L^{'} > 0$ (in places of G, δ and L in Theorem 1.6.9 of [G5]). Apply Lemma 2.7 to the set $\widetilde{E} = E_1 \cup E_2 \cup F_1$ and $\widetilde{\delta} = \frac{1}{3}\min(\varepsilon, \delta, \delta_1, \delta_2)$ to obtain the diagram



with $\beta^{'}$ being $\widetilde{E} - \widetilde{\delta}$ multiplicative and

$$\| \ \iota \circ \beta^{'}(f) - \phi^{'}(f) \ \| < \widetilde{\delta}, \ \forall \ f \in \widetilde{E}.$$

Let $L = L^{'} \cdot rank(\mathbf{1}_A)$, and let $\beta_1 : A \longrightarrow M_L(B^{'})$ be any unital homomorphism defined by point evaluation. Then by Theorem 1.6.9 of [G5], there is a unitary $u \in M_{L+1}(B)$ such that

$$\|u((\beta^{'} \oplus \beta_1) \circ \alpha^{'}(f))u^* - (\beta^{'} \oplus \beta_1) \circ \alpha_1(f)\| < 8\varepsilon_1, \ \forall \ f \in \widetilde{G}_1^i.$$

By the choice of E_1, there is a homomorphism

$$\widetilde{\xi} : C \longrightarrow M_{L+1}(B^{'}),$$

such that

$$\|\widetilde{\xi}(f) - u((\beta^{'} \oplus \beta) \circ \xi_2(f))u^*\| < \varepsilon_1, \ \forall \ f \in \xi_1(\widetilde{G}_1^i).$$

Define $B_2 = M_{L+1}(B^{'}), K = L_1(L+1), A^{'} = M_K(A) = M_{L+1}(M_{L_1}(A))$,

$$\psi : B_1^i \longrightarrow B_2 \ \text{ by } \ \psi = \widetilde{\xi} \circ \xi_1 : B_1^i \xrightarrow{\xi_1} C \xrightarrow{\widetilde{\xi}} B_2,$$

$$\beta : A \longrightarrow M_{L+1}(B^{'}) \ \text{ by } \ \beta = \beta^{'} \oplus \beta_1,$$

and

$$\phi : A \longrightarrow M_{L+1}(M_{L_1}(A)) \ \text{ by } \ \phi = \phi^{'} \oplus ((\iota \otimes id_L) \circ \beta_1),$$

(note that β_1 is a homomorphism) to finish the proof.

□

2.13. Recall for $A = \bigoplus_{i=1}^{t} M_{k_i}(C(X_i))$, where X_i are path connected simplicial complexs, we use the notation $r(A)$ to denote $\bigoplus_{i=1}^{t} M_{k_i}(\mathbb{C})$ which could be considered to be the subalgebra consisting of all t-tuples of constant function from X_i to $M_{k_i}(\mathbb{C})$ $(i = 1, 2, \cdots, t)$. Fixed a base point $x_i^0 \in X_i$ for each X_i, one defines a map $r : A \longrightarrow r(A)$ by

$$r(f_1, f_2, \cdots, f_t) = (f_1(x_1^0), f_2(x_2^0), \cdots, f_t(x_t^0)) \in r(A).$$

We have the following Corollary

Corollary 2.14. Let $B_1 = \oplus B_1^j$, where B_1^j are either of form $M_{k(j)}(C(X_j))$, with X_j being one of $\{pt\}$, [0,1], S^1, $\{T_{II,k}\}_{k=2}^{\infty}$ or one of form $M_{k(j)}(I_{l(j)})$. Let $\alpha_1; B_1 \longrightarrow A$ be a homomorphism, where A is a direct sum of matrix algebras over $\{pt\}, [0,1], S^1, \{T_{II,k}\}_{k=2}^{\infty}, \{T_{III,k}\}_{k=2}^{\infty}$ and S^2. Let $\varepsilon_1 > 0$ and let $\widetilde{E}(= \oplus \widetilde{E}^i) \subset E(= \oplus E^i) \subset B_1(= \oplus B_1^i)$ be two finite subset with the condition

$$\omega(\widetilde{E}^i) < \varepsilon_1, \text{ if } B_1^i = M_{\bullet}(C(Y_i)) \text{ with } Y_i \in \{T_{II,k}\}_{k=2}^{\infty}.$$

Let $F \subset A$ be any finite set, $\varepsilon_2 > 0$, $\delta > 0$. Then there exists a diagram



where $A^{'} = M_L(A)$, and B_2 is a direct sum of matrix algebras over space: $\{pt\}$, [0,1], S^1, $\{T_{II,k}\}_{k=2}^{\infty}$ and dimension drop algebras, with the following properties.
(1) ψ is a homomorphism, α_2 is a injective homomorphism and ϕ is a unital simple embedding.

(2) $\beta \in Map(A, B_2)_1$ is $F_1 - \delta$ multiplicative.
(3) for $g \in \widetilde{E}^i$ with $B_1^i = M_\bullet(C(X_i))$, $X_i \in \{T_{II,k}\}_{k=2}^\infty$ we have

$$\|(\beta \oplus r)(g) - (\psi \oplus (r \circ \alpha_1))(g)\| \leqslant 10\varepsilon_1;$$

and for $g \in E^i(\supset \widetilde{E}^i)$ where B_1^i is not of form $M_\bullet(C(T_{II,k}))$, we have

$$\|(\beta \oplus r)(g) - (\psi \oplus (r \circ \alpha_1))(g)\| < \varepsilon_1;$$

and for $f \in F$

$$\|(\alpha_2 \oplus id) \circ (\beta \oplus r)(f) - (\phi \oplus r)(f)\| < \varepsilon_1.$$

(4) For B_2^i of form $M_\bullet(C(T_{II,k}))$,

$$\omega(\pi_i(\beta(F) \cup \psi(E))) < \varepsilon_2,$$

where π_i is the canonical projection from B_2 to B_2^i.

Remark: In the application of the above Corollary, we will denote the map $\beta \oplus r$ by β and $\psi \oplus (r \circ \alpha_1)$ by ψ.

§3. The proof of main theorem

In this section, we prove the following main theorem.

Theorem 3.1. Suppose $\lim(A_n = \bigoplus_{i=1}^{t_n} M_{[n,i]}(C(X_{n,i})), \phi_{n,m})$ is an AH inductive limit with $X_{n,i}$ being among the spaces $\{pt\}, [0,1], S^1, \{T_{II,k}\}_{k=2}^\infty, \{T_{III,k}\}_{k=2}^\infty$, such that the limit algebra A has ideal property. Then there is another inductive system $(B_n = \oplus B_n^i, \psi_{n,m})$ with same limit algebra, where B_n^i are either $M_{[n,i]'}(C(Y_{n,i}))$ with $Y_{n,i}$ being one of $\{pt\}, [0,1], S^1, \{T_{II,k}\}_{k=2}^\infty$ (but without $T_{III,k}$ and S^2), or dimension drop algebra $M_{[n,i]'}(I_{k(n,i)})$.

Proof. Let $\varepsilon_1 > \varepsilon_2 > \varepsilon_3 > \cdots$ be a sequence of positive number with $\sum \varepsilon_n < +\infty$. We need to construct the intertwining diagram



satisfying the following conditions
(0.1) $(A_{s(n)}, \phi_{s(n),s(m)})$ is a sub-inductive system of $(A_n, \phi_{n,m}), (B_n, \psi_{n,m})$ is an inductive system of direct sum of matrix algebras over the spaces $\{pt\}, [0,1], S^1, T_{II,k}$ and dimension drop algebra $M_\bullet(I_{k(n,i)})$.
(0.2) Choose $\{a_{i,j}\}_{j=1}^\infty \subset A_{s(i)}$ and $\{b_{i,j}\}_{j=1}^\infty \subset B_i$ to be countable dense subsets of unit balls of $A_{s(i)}$ and B_i, respectively. F_n are subsets of unit balls of $A_{s(n)}$, and $\widetilde{E_n} \subset E_n$ are both subsets of unit balls of B_n satisfying

$$\phi_{s(n),s(n+1)}(F_n) \cup \alpha_{n+1}(E_{n+1}) \cup \bigcup_{i=1}^{n+1} \phi_{s(i),s(n+1)}(\{a_{i1}, a_{i2}, \cdots, a_{in+1}\}) \subset F_{n+1},$$

$$\psi_{n,n+1}(E_n) \cup \beta_n(F_n) \subset \widetilde{E}_{n+1} \subset E_{n+1},$$

and

$$\bigcup_{i=1}^{n+1} \psi_{i,n+1}(\{b_{i1}, b_{i2}, \cdots, b_{in+1}\}) \subset E_{n+1}.$$

(Here $\phi_{n,n} : A_n \longrightarrow A_n$, and $\psi_{n,n} : B_n \longrightarrow B_n$ are understand as identity maps.)
(0.3) β_n are $F_n - 2\varepsilon_n$ multiplicative and α_n are homomorphism.
(0.4) for all $g \in \widetilde{E}_n$,

$$\|\psi_{n,n+1}(g) - \beta_n \circ \alpha_n(g)\| < 14\varepsilon_n,$$

and for all $f \in F_n$,

$$\|\phi_{s(n),s(n+1)}(f) - \alpha_{n+1} \circ \beta_n(f)\| < 14\varepsilon_n.$$

(0.5) For any block B_n^i with spectrum $T_{II,k}$, we have $\omega(\widetilde{E}_n^i) < \varepsilon_n$, where $\widetilde{E}_n^i = \pi_i(\widetilde{E}_n)$ for $\pi_i : B_n \longrightarrow B_n^i$ the canonical projections.

The diagram will be constructed inductively. First, let $B_1 = \{0\}, A_{s(1)} = A_1, \alpha_1 = 0$. Let $b_{1j} = 0 \in B_1$ for $j = 1, 2, ...$ and let $\{a_{1j}\}_{j=1}^\infty$ be a countable dense subset of the unit ball of $A_{s(1)}$. And let $\widetilde{E}_1 = E_1 = \{b_{11}\} = B_1$ and $F_1 = \bigoplus_{i=1}^{t_1} F_1^i$, where $F_1^i = \pi_i(\{a_{11}\}) \subset A_1^i$.

As inductive assumption, assume that we already have the diagram



and for each $i = 1, 2, \cdots, n$, we have dense subsets $\{a_{ij}\}_{j=1}^{\infty} \subset$ the unit ball of $A_{s(i)}$ and $\{b_{ij}\}_{j=1}^{\infty} \subset$ the unit ball of B_i satisfying the conditions (0.1)-(0.5) above. We have to construct the next piece of the diagram


$F_n \subset A_{s(n)}$ $\phi_{s(n),s(n+1)}$ $A_{s(n+1)} \supset F_{n+1}$
α_n β_n α_{n+1}
$\widetilde{E}_n \subset E_n \subset B_n$ $B_{n+1} \supset E_{n+1} \supset \widetilde{E}_{n+1}$
$\psi_{n,n+1}$

to satisfy the condition (0.1)-(0.5).

Among the conditions for induction assumption, we will only use the conditions that α_n is a homomorphism and (0.5) above.
Step 1. We enlarge $\widetilde{E}_n$ to $\bigoplus_i \pi_i(\widetilde{E}_n^i)$ and E_n to $\bigoplus_i \pi_i(E_n)$. Then $\widetilde{E}_n(= \oplus \widetilde{E}_n^i) \subset E_n(= \oplus E_n)$ and for each B_n^i with spectrum $T_{II,k}$, we have $\omega(E_n^i) < \varepsilon_n$ from induction assumption (0.5). By proposition 2.9 applied to $\alpha_n : B_n \to A_{s(n)}, \widetilde{E}_n \subset E_n \subset B_n, F_n \subset A_{s(n)}$ and $\varepsilon_n > 0$, there are $A_{m_1}(m_1 > s(n))$, two othogonal projections $P_0, P_1 \in A_{m_1}$ with $\phi_{s(n),m_1}(\mathbf{1}_{A_{s(n)}}) = P_0 + P_1$ and P_0 trivial, a C^*-algebra C—a direct sum of matrix algebras over $C[0,1]$ or $\mathbb{C}$, a unital map $\theta \in Map(A_{s(n)}, P_0 A_{m_1} P_0)_1$, a unital homomorphism $\xi_1 \in Hom(A_{s(n)}, C)_1$, a unital homomorphism $\xi_2 \in Hom(C, P_1 A_{m_1} P_1)_1$ such that
(1.1) $\|\phi_{s(n),m_1}(f) - \theta(f) \oplus (\xi_2 \circ \xi_1)(f)\| < \varepsilon_n$ for all $f \in F_n$.
(1.2) θ is $F_n - \varepsilon$ multiplicative and $F := \theta(F_n)$ satisfy $\omega(F) < \varepsilon_n$.
(1.3) $\|\alpha(g) - \theta \circ \alpha_n(g)\| < 3\varepsilon_n$ for all $g \in \widetilde{E}_n$.

Let all the blocks of C be parts of C^*-algebra B_{n+1}. That is

$$B_{n+1} = C\oplus \text{ (some other blocks)}.$$

The map $\beta_n : A_{s(n)} \to B_{n+1}$, and the homomorphism $\psi_{n,n+1} : B_n \to B_{n+1}$ are defined by $\beta_n = \xi_1 : A_{s(n)} \to C(\subset B_{n+1})$ and $\psi_{n,n+1} = \xi_1 \circ \alpha_n : B_n \to C(\subset B_{n+1})$ for the blocks of $C(\subset B_{n+1})$. For this part, β_n is also a homomorphism.
Step 2. Let $A = P_0 A_{m_1} P_0, F = \theta(F_n)$. Since P_0 is a trivial projection,

$$A \cong \oplus M_{l_i}(C(X_{m_1,i})).$$

Let $r(A) := \oplus M_{l_i}(\mathbb{C})$ and $r : A \to r(A)$ be as in 2.13. Applying corollary 2.14 and its remark to $\alpha : B_n \to A, \widetilde{E}_n \subset E_n \subset B_n$ and $F \subset A$, we obtain the following diagram



such that
(2.1) B is a direct sum of matrix algebras over $\{pt\}, [0,1], S^1, T_{II,k}$ and dimension drop algebra.
(2.2) $\alpha^{'}$ is an injective homomorphism and β is $F - \varepsilon_n$ multiplicative.
(2.3) $\phi : A \to M_L(A)$ is a unital simple embedding and $r : A \to r(A)$ is as in 2.13.
(2.4) $\|\beta \circ \alpha(g) - \psi(g)\| < 10\varepsilon_n$ for all $g \in \widetilde{E}_n$ and $\|(\phi \oplus r)(f) - \alpha^{'} \circ \beta(f)\| < \varepsilon_n$ for all $f \in F(:= \theta(F_n))$.
(2.5) $\omega(\pi_i(\psi(E_n)) \cup \beta(F)) < \varepsilon_{n+1}$ (note that $\beta(F) = \beta \circ \theta(F_n)$), for B_n^i being of form $M_\bullet(C(X))$ with $X \in \{T_{II,k}\}_{k=2}^{\infty}$.

Let all the blocks B be also part of B_{n+1}, that is

$$B_{n+1} = C \oplus B\oplus \text{ (some other blocks)}.$$

The maps $\beta_n : A_{s(n)} \longrightarrow B_{n+1}, \psi_{n,n+1} : B_n \longrightarrow B_{n+1}$ are defined by

$$\beta_n := \beta \circ \theta : A_{s(n)} \xrightarrow{\theta} A \xrightarrow{\beta} B(\subset B_{n+1}),$$

and

$$\psi_{n,n+1} := \psi : B_n \to B(\subset B_{n+1}),$$

for the blocks of $B(\subset B_{n+1})$. This part of β_n is $F_n - 2\varepsilon_n$ multiplicative, since θ is $F_n - \varepsilon_n$ multiplicative, β is $F - \varepsilon_n$ multiplicative and $F = \theta(F_n)$.
Step 3. By Lemma 3.15 of [GJLP2] applied to $\phi \oplus r : A \to M_L(A) \oplus r(A)$, there is an A_{m_2} and there is a unital homomorphism

$$\lambda : M_L(A) \oplus r(A) \to RA_{m_2}R,$$

where $R = \phi_{m_1,m_2}(P_0)$ (write R as $\bigoplus_j R^j \in \bigoplus_j A_m^j$) such that the diagram



satisfies the following conditions:
(3.1) $\lambda \circ (\phi \oplus r)$ is homotopy equivalent to

$$\phi^{'} := \phi_{m_1,m_2}|_A.$$

Step 4. Applying Theorem 1.6.9 of [G5] to finite set $F \subset A$ (with $\omega(F) < \varepsilon_n$) and to two homotopic homomorphisms $\phi^{'}$ and $\lambda \circ (\phi \oplus r) : A \to RA_{m_2}R$ (with $RA_{m_2}R$ in place of C in Theorem 1.6.9 of [G5]), we obtain a finite set $F^{'} \subset RA_{m_2}R$, $\delta > 0$ and $L > 0$ as in the Theorem.

Let $G = \oplus\pi_i(\psi(E_n) \cup \beta(F)) = \oplus G^i$. Then by (2.5), we have $\omega(G^i) < \varepsilon_{n+1}$, if B^i is of form $M_\bullet(C(T_{II,k}))$. By Proposition 2.8 applied to $RA_{m_2}R$ and

$$\lambda \circ \alpha^{'} : B \to RA_{m_2}R,$$

finite set $G \subset B$, $F^{'} \cup (\phi_{m_1m_2}\ |_A\ (F)) \in RA_{m_2}R$, $min(\varepsilon_n, \delta) > 0$ (in place of ε) and $L > 0$, there are $A_{s(n+1)}$, mutually orthogonal projections $Q_0, Q_1, Q_2 \in A_{s(n+1)}$ with $\phi_{m_2,s(n+1)}(R) = Q_0 \oplus Q_1 \oplus Q_2$, a C^*-algebra D—a direct sum of matrix algebras over C[0,1] or $\mathbb{C}$—, a unital map $\theta_0 \in \mathrm{Map}(RA_{m_2}R, Q_0A_{s(n+1)}Q_0)$, and four unital homomorphisms

$$\theta_1 \in Hom(RA_{m_2}R, Q_1A_{s(n+1)}Q_1)_1, \xi_3 \in Hom(RA_{m_2}R, D)_1, \xi_4 \in Hom(D, Q_2A_{s(n+1)}Q_2)_1$$

and $\alpha^{''} \in Hom(B, (Q_0+Q_1)A_{s(n+1)}(Q_0+Q_1))_1$ such that the following is true.
(4.1) $\|\phi_{m_2,s(n+1)}(f) - ((\theta_0+\theta_1) \oplus \xi_4 \circ \xi_3)(f)\| < \varepsilon_n$, for all $f \in \phi_{m_1,m_2}|_A(F) \subset RA_{m_2}R$.
(4.2) $\|\alpha^{''}(g) - (\theta_0+\theta_1) \circ \lambda \circ \alpha^{'}(g)\| < 3\varepsilon_{n+1} < 3\varepsilon_n$, $\forall\ g \in G$.
(4.3) θ_0 is $F^{'} - min(\varepsilon_n, \delta)$ multiplicative and θ_1 satisfies that

$$\theta_1^{i,j}([q]) > L \cdot [\theta_0^{i,j}(R^i)],$$

for any non zero projection $q \in R^iA_{m_1}R^i$.
By Theorem 1.6.9 of [G5], there is a unitary $u \in (Q_0 \oplus Q_1)A_{s(n+1)}(Q_0+Q_1)$ such that

$$\|(\theta_0+\theta_1) \circ \phi^{'}(f) - Adu \circ (\theta_0+\theta_1) \circ \lambda \circ (\phi \oplus r)(f)\| < 8\varepsilon_n,$$

for all $f \in F$.
Combining with second inequality of (2.4), we have

(4.4)$\|(\theta_0+\theta_1)\circ\phi^{'}(f)-Adu\circ(\theta_0+\theta_1)\circ\lambda\circ\alpha^{'}\circ\beta(f)\|<9\varepsilon_n$ for all $f\in F$.
Step 5. Finally let all blocks of D be the rest of B_{n+1}. Namely, let

$$B_{n+1}=C\oplus B\oplus D,$$

where C is from Step 1, B is from Step 2 and D is from Step 4.

We already have the definition of $\beta_n:A_{s(n)}\to B_{n+1}$ and $\psi_{n,n+1}:B_n\to B_{n+1}$ for those blocks of $C\oplus B\subset B_{n+1}$ (from Step 1 and Step 2). The definition of β_n and $\psi_{n,n+1}$ for blocks of D and the homomorphism $\alpha_{n+1}:C\oplus B\oplus D\to A_{s(n+1)}$ will be given below.

The part of $\beta_n:A_{s(n)}\to D(\subset B_{n+1})$ is defined by

$$\beta_n=\xi_3\circ\phi^{'}\circ\theta:A_{s(n)}\xrightarrow{\theta}A\xrightarrow{\phi}RA_{m_2}R\xrightarrow{\xi_3}D.$$

(Recall that $A=P_0A_{m_2}P_0$ and $\phi^{'}=\phi_{m_1,m_2}|_A$.) Since θ is $F_n-\varepsilon_n$ multiplicative, and $\phi^{'}$ and ξ_3 are homomorphism, we know this part of β_n is $F_n-\varepsilon_n$ multiplicative.

The part of $\psi_{n,n+1}:B_n\to D(\subset B_{n+1})$ is defined by

$$\psi_{n,n+1}=\xi_3\circ\phi^{'}\circ\alpha:B_n\xrightarrow{\alpha}A\xrightarrow{\phi^{'}}RA_mR\xrightarrow{\xi_3}D,$$

which is a homomorphism.

The homomorphism $\alpha_{n+1}:C\oplus B\oplus D\to A_{s(n+1)}$ is defined as following.

Let $\phi^{''}=\phi_{m_1,s(n+1)}|_{P_1A_{m_1}P_1}:P_1A_{m_1}P_1\longrightarrow\phi_{m_1,s(n+1)}(P_1)A_{s(n+1)}\phi_{m_1,s(n+1)}(P_1)$, where P_1 is from Step 1. Define

$$\alpha_{n+1}|_C=\phi^{''}\circ\xi_2:C\xrightarrow{\xi_2}P_1A_{m_1}P_1\xrightarrow{\phi^{''}}\phi_{m_1,s(n+1)}(P_1)A_{s(n+1)}\phi_{m_1,s(n+1)}(P_1),$$

where ξ_2 is from Step 1.

$$\alpha_{n+1}|_B=Adu\circ\alpha^{''}:B\xrightarrow{\alpha^{''}}(Q_0\oplus Q_1)A_{s(n+1)}(Q_0+Q_1)\xrightarrow{Adu}(Q_0\oplus Q_1)A_{s(n+1)}(Q_0+Q_1)$$

where $\alpha^{''}$ is from Step 4, and define

$$\alpha_{n+1}|_D=\xi_4:D\to Q_2A_{s(n+1)}Q_2.$$

Finally choose $\{a_{n+1,j}\}_{j=1}^{\infty}\subset A_{s(n+1)}$ and $\{b_{n+1,j}\}_{j=1}^{\infty}\subset B_{n+1}$ to be countable dense subsets of the unit balls of $A_{s(n+1)}$ and B_{n+1}, respectively. And choose

$$F_{n+1}^{'}=\phi_{s(n),s(n+1)}(F_n)\cup\alpha_{n+1}(E_{n+1})\cup\bigcup_{i=1}^{n+1}\phi_{s(i),s(n+1)}(\{a_{i1},a_{i2},\cdots,a_{in+1}\}),$$

$$E_{n+1}^{'} = \psi_{n,n+1}(E_n) \cup \beta_n(F_n) \cup \bigcup_{i=1}^{n+1} \psi_{i,n+1}(\{b_{i1}, b_{i2}, \cdots, b_{in+1}\}),$$

$$\widetilde{E}_{n+1}^{'} = \psi_{n,n+1}(E_n) \cup \beta_n(F_n) \subset E_{n+1}^{'}.$$

Define $F_{n+1}^i = \pi_i(F_{n+1}^{'})$ and $F_{n+1} = \bigoplus_i F_{n+1}^i$, $E_{n+1}^i = \pi_i(E_{n+1}^{'})$ and $E_{n+1} = \oplus_i E_{n+1}^i$. For those block B_{n+1}^i inside the algebra B define $\widetilde{E}_{n+1}^i = \pi_i(\widetilde{E}_{n+1})$. For those blocks inside C and D, define $\widetilde{E}_{n+1}^i = E_{n+1}^i$. And finally let $E_{n+1} = \bigoplus_i \widetilde{E}_{n+1}^i$. Note all the blocks with spectrum $T_{II,k}$ are in B. And (2.5) tells us that for those blocks $\omega(\widetilde{E}_{n+1}^i) < \varepsilon_{n+1}$. Thus we obtain the following diagram



Step 6.Now we need to verify all the condition (0.1)-(0.5) for the above diagram.

From the end of Step 5, we know (0.5) holds, (0.1)-(0.2) hold from the construction (see the construction of B, C, D in Step 1, 2 and 4, and $\widetilde{E}_{n+1} \subset E_{n+1}, F_{n+1}$ is the end of Step 5). (0.3) follows from the end of Step 1, the end of Step 2 and the part of definition of β_n for D from Step 5.

So we only need to verify (0.4).

Combining (1.1) with (4.1), we have

$$\|\phi_{s(n),s(n+1)}(f) - [(\phi^{''} \circ \xi_2 \circ \xi_1) \oplus (\theta_0 + \theta_1) \circ \phi^{'} \circ \theta \oplus (\xi_4 \circ \xi_3 \circ \phi^{'} \circ \theta)(f)](f)\| < \varepsilon_n + \varepsilon_n = 2\varepsilon_n$$

for all $f \in F_n$ (recall that $\phi^{''} = \phi_{m_1,s(n+1)}|_{P_1 A_{m_1} P_1}, \phi^{'} := \phi_{m_1,m_2}|_{P_0 A_{m_1} P_o}$).

Combining with (4.2) and (4.4), and definition of β_n and α_{n+1}, the above inequality yields

$$\|\phi_{s(n),s(n+1)}(f) - (\alpha_{n+1} \circ \beta_{n+1})(f)\| < 9\varepsilon_n + 3\varepsilon_n + 2\varepsilon_n = 14\varepsilon_n, \ \forall \ f \in F_n.$$

Combining (1.3), the first inequality of (2.4) and the definition of β_n and $\psi_{n,n+1}$, we have

$$\|\psi_{n,n+1}(g) - (\beta_n \circ \alpha_n)(g)\| < 10\varepsilon_n + 3\varepsilon_n < 14\varepsilon_n, \ \forall \ g \in \widetilde{E}_n.$$

So we obtain(0.4). The theorem follows from Proposition 4.1 of [GJLP2].

□

Note that if $q \in M_l(I_k)$, then $qM_k(I_k)q$ isomorphic to $M_{l_1}(I_k)$. Combining with the main theorem of [GJLP2] (see Theorem 4.2, and 2.7 of [GJLP2]) we have

Theorem 3.2. Suppose that $A = \lim(A_n = \oplus P_{n,i} M_{[n,i]}(C(X_{n,i}))P_{n,i})$ is an AH inductive limit with $dim(X_{n,i}) \leqslant M$ for a fixed positive integer M such that limit algebra A has ideal property. Then A can be rewrite as inductive limit $\lim(B_n = \oplus B_n^i, \psi_{n,m})$, where either $B_n^i = Q_{n,i} M_{[n,i]'}(C(Y_{n,i}))Q_{n,i}$ with $Y_{n,i}$ being one of the spaces $\{pt\}, [0,1], S^1, \{T_{II,k}\}_{k=2}^{\infty}$ or $B_n^i = M_{[n,i]'}(I_{l_{(n,i)}})$ a dimension drop algebra.

References

[Blackadar] B. Blackadar, Matricial and ultra-matricial topology, Operator Algebras, Mathematical Physics, and Low Dimensional Topology (R. H. Herman and B Tanbay eds) A K Peter, Massachusetts (1993), 11-38

[D1] M. Dadarlat, Approximately unitarily equiralent, morphisms and inductive limit C^*-algebras, K-theory 9(1995), 117-137

[D2] M. Dadarlat, Reduction to dimesion these of local spectra of Real rank zero C^*-algebras, J. Reine Angew. Math. 460(1995) 189-212

[DG] M. Dadarlat and G. Gong, A classification result for approximately homogeneous C^*-algebras of real rank zero, Geometric and Functional Analysis, 7(1997) 646-711

[DN] M. Dadarlat and A. Nemethi, Sharp theory and (connective) K-theory, J. Operator Theory 23(1990) 207-291

[Ell1] G. A. Elliott, On the classification of C^*-algebras of real rank zero, J. Reine Angew. Math. 443(1993) 263-290

[Ell2] G. A. Elliott, A classification of certain simple C^*-algebras, Quantum and Non-Commutative Analysis, Kluwer, Dordrecht, (1993), pp, 373-388

[Ell3] G. A. Elliott, A classification of certain simple C^*-algebras, II, J. Ramaunjan Math. Soc. 12 (1997), 97-134

[EG1] G. A. Elliott and G. Gong, On the inductive limits of matrix algebras over two-tori, American. J. Math 118(1996) 263-290

[EG2] G. A. Elliott and G. Gong, On the classification of C^*-algebras of real rank zero, II. Ann. of Math 144(1996) 497-610

[EGL1] G. A. Elliott, G. Gong and L. Li, On the classification of simple inductive limit C^*-algebras, II; The isomorphism Theorem, Invent. Math. 168(2)(2007) 249-320

[EGL2] G. A. Elliott, G. Gong and L. Li, Injectivity of the connecting maps in AH inductive limit systems, Canand. Math. Bull. 26(2004) 4-10

[G1] G. Gong, Approximation by dimension drop C^*-algebras and classification, C. R. Math. Rep. Acad. Sci Can. 16(1994)40-44

[G2] G. Gong, Classification of C^*-algebras of real rank zero and unsuspended E-equivalent types, J. Funct. Anal. 152(1998) 281-329

[G3-4] G. Gong, On inductive limit of matrix algebras over higher dimension spaces, Part I, II, Math Scand. 80(1997) 45-60, 61-100

[G5] G. Gong, On the classification of simple inductive limit C^*-algebras, I: Reduction Theorems. Doc. Math. 7(2002) 255-461

[GJL] G. Gong, C. Jiang, and L. Li, A classification of inductive limit C^*-algebras with ideal property, arXiv:1607.07581

[GJLP1] G. Gong, C. Jiang, L. Li, C. Pasnicu, AT structure of AH algebras with ideal property and torsion free K-theory, J. Func. Anal. 58(2010) 2119-2143

[GJLP2] G. Gong, C. Jiang, L. Li, C. Pasnicu, A Reduction theorem for AH algebras with ideal property, arXiv:1607.07575

[Ji-Jiang] K. Ji and C. Jiang, A complete classification of AI algebra with ideal property, Canadian. J. Math, 63(2), (2011), 381-412

[Jiang] C. Jiang, A classification of non simple C^*-algebras of tracial rank one:Inductive limit of finite direct sums of simple TAI C^*-algebras, J. Topol. Anal. 3 No.3(2011), 385-404

[Li1] L. Li, On the classification of simple C^*-algebras: Inductive limit of matrix algebras tree, Men Amer. Math, Soc. 127(605)(1997)

[Li2] L. Li, Simple inductive limit C^*-algebras: Spectra and approximation by interral algebras, J. Reine Angew Math 507(1999) 57-79

[Li3] L. Li, Classification of simple C^*-algebras: Inductive limit of matrix algebras over 1-dimensional spaces, J. Func. Anal. 192(2002) 1-51

[Li4] L. Li, Reduction to dimension two of local spectrum for simple AH algebras, J. of Ramanujia Math. Soc. 21 No.4(2006) 365-390

[Pasnicu1] C. Pasnicu, On inductive limit of certain C^*-algebras of the form $C(x)\otimes F$, Trans. Amer. Math. Soc. 310(2)(1988) 703-714

[Pasnicu2] C. Pasnicu, Shape equiralence, nonstable K-theory and AH algebras, Pacific J. Math 192(2000) 159-182